

Pressemitteilung

Julius-Maximilians-Universität Würzburg

Robert Emmerich

03.03.2022

<http://idw-online.de/de/news789438>

Forschungsergebnisse, Wissenschaftliche Publikationen
Biologie
überregional



What Makes Plants Electrically Excitable

In humans, only nerves and muscle cells are electrically excitable, whereas in plants almost all cells are. This is due to a sophisticated mechanism in an ion channel of the vacuole.

Plant cells use electrical signals to process and transmit information. In 1987, as a postdoc of Erwin Neher in Göttingen, biophysicist Rainer Hedrich discovered an ion channel in the central vacuole of the plant cell, which is activated by calcium and electrical voltage, using the patch-clamp technique (Nobel Prize for Neher and Sakmann 1991).

In 2019, Hedrich's team at Julius-Maximilians-Universität Würzburg (JMU) identified this TPC₁ channel as an important element for electrical communication in plants. If the channel fails, signal transmission is slowed down. If it is hyperactive, i.e. open for too long, the plant is highly stressed and has problems growing.

These reactions make it clear: plants must strictly control the opening time of the ion channel TPC₁ so that the electrical communication between their cells runs smoothly.

The structure explains how the channel is switched on

A publication in the journal PNAS (Proceedings of the National Academy of Sciences of the USA) now provides new insights into the molecular function and regulation of the TPC₁ channel. This was achieved by combining two areas of expertise: A JMU team led by Rainer Hedrich and Irene Marten was responsible for plant biophysics, and a group led by Robert M. Stroud and Sasha Dickinson from the University of California in San Francisco was responsible for structural biology.

Using high-resolution cryo-electron microscopic imaging, the US team clearly demonstrates that massive conformational changes occur in several protein domains before the channel opens. An electrical stimulus initiates a rotating movement of the voltage sensor domain. This flaps away amino acid residues that serve as binding sites for inhibitory calcium ions in the vacuolar entry region of the channel - clearing the way for ion flow.

The vacuolar calcium content keeps the voltage sensor in check

Hedrich and Marten's team was able to show that the TPC₁ channel is switched on when the calcium level in the cell plasma rises in response to external stimuli. An increase in the calcium level in the vacuole, on the other hand, slows down an excessive ion flux through the channel and virtually renders the vacuolar membrane insensitive to Calcium-dependent electrical stimuli.

The newly discovered binding site for vacuolar calcium ions in the ion-conducting pore of the channel plays a crucial role in this process.

"We were able to elucidate the function of this channel domain with patch clamp measurements," says JMU professor Irene Marten. "When calcium binds to the vacuolar pore-binding site, negative feedback with the voltage sensor occurs, which means that the movement of the voltage sensor domain is strongly impaired. As a result, the channel remains closed and there is no electrical excitation of the vacuole. If, on the other hand, the pore amino acid residues are removed from the ion transport pathway, no vacuolar calcium binding occurs and the channel opening is strongly facilitated".

Question about the evolution of TPC₁ channels

The publication in PNAS further contributes to the fact that the plant ion channel TPC₁ is now one of the best understood voltage-dependent ion channels. This knowledge can help to better understand TPC₁-dependent processes in animal cells as well.

What will the scientists do next? "We are looking into the question of whether the TPC₁ channels of different plant species differ in terms of regulation and also in other properties, and whether this opens up new possibilities for adaptation to the environment," says Rainer Hedrich. "In doing so, we are also taking into account regulators that play a role in animal TPC₁ channels. The studies should also give us an insight into the evolution of TPC₁ channels."


wissenschaftliche Ansprechpartner:

Prof. Dr. Irene Marten, Chair of Botany I (Molecular Plant Physiology and Biophysics), University of Würzburg, T +49 931 31-86118, marten@botanik.uni-wuerzburg.de

Prof. Dr. Rainer Hedrich, Chair of Botany I (Molecular Plant Physiology and Biophysics), University of Würzburg, T +49 931 31-86100, hedrich@botanik.uni-wuerzburg.de

Originalpublikation:

Molecular basis of multistep voltage activation in plant two-pore channel 1. PNAS, 24.02.2022, 119 (9) e2110936119 | Free Access: <https://doi.org/10.1073/pnas.2110936119>



3-Dimensional sketch of the TPC₁ channel protein looking at the vacuolar pore entrance from above. Positional changes of amino acid residues such as E605 during the transition from a closed channel state to a partially open pore state.

Thomas Müller

University of Würzburg