(idw)

Press release

Klinikum der Universität München Philipp Kressirer

10/20/2021 http://idw-online.de/en/news777826 Research projects, Scientific Publications Medicine transregional, national

LMU KLINIKUM

Spread of SARS-CoV-2 in Ethiopia is underestimated

The Journal "The Lancet Global Health" has published the results of a longitudinal cohort study on the spread of SARS-CoV-2 in Ethiopia. In an Ethiopian-German research collaboration, the Division of Infectious Diseases and Tropical Medicine at the LMU University Hospital investigated blood samples of frontline healthcare workers and residents from urban and rural communities for antibodies. The results suggest that the true COVID-19 prevalence is much higher than previously reported official figures. Therefore, the research team recommends a realignment of the vaccination strategy for Africa.

Since the first reported COVID-19 case in Ethiopia in March 2020, the actual spread and burden of SARS-CoV-2 in the country and in Africa has remained largely unknown. Thus far, official data have provided little information in this regard. Contrary to fears of a humanitarian crisis, relatively few COVID-19 cases (of death) were reported for the African continent at the onset of the pandemic. However, recent studies indicate a high prevalence of SARS-CoV-2 among health workers (e.g., 41.2 percent in the Democratic Republic of the Congo, 45.1 percent in Nigeria). Other sporadic reports provide figures as high as 60 percent among blood donors in South Africa. Therefore, the question arises as to how severely the African continent is affected by the virus and what this means for effective pandemic control strategies in Africa.

In an international collaboration of the Division of Infectious Diseases and Tropical Medicine at the LMU University Hospital (Dr. Arne Kroidl and Dr. Andreas Wieser) and of the Ethiopian partners of Jimma University (Jimma Medical Center, JMC, Professor Esayas Kebede Gudina, PhD) and the St. Paul's Hospital in Addis Abeba (SPHMMC, Solomon Ali, PhD), together with the Helmholtz Zentrum Munich, the LMU, the Technical University Munich and the University of Bonn, the research team* conducted a study on the spread of SARS-CoV-2 by antibody detection. In their research project, they analysed blood samples from frontline healthcare workers, urban as well as rural community residents in Jimma and Addis Ababa.

The aim of the cohort study was to obtain epidemiological data on seroprevalence (frequency of specific antibodies in blood serum) and seroincidence (increase in the proportion of antibody-positive individuals) over time in Africa for the first time.

The research was funded by the Bavarian State Chancellery. Bavaria's State Minister for European and International Affairs, Melanie Huml, on the research project: "One thing is clear: we can only overcome global challenges such as the COVID-19 pandemic collectively. That is why Bavaria has supported the Bavarian-Ethiopian research on the dissemination of COVID-19 antibodies with 500,000 euros. The research results demonstrate the value of this investment. Based on these results, the containment of COVID-19 on the African continent can be achieved more effectively. I am pleased that the long-standing partnership between Bavarian and Ethiopian stakeholders in the health sector benefits all partners."

In the period between August 2020 and April 2021, the researchers conducted serological testing of blood samples from healthcare professionals at two teaching hospitals in Jimma city and surrounding rural communities, as well as in the Ethiopian capital Addis Ababa. This was done in a total of three study rounds.

(idw)

Dramatic increase in SARS-CoV-2 seroprevalence over time:

The results showed that SARS-CoV-2 seroprevalence among medical staff had increased dramatically during the study period: at the Addis Ababa hospital, the team recorded an increase from 10.9 percent in August 2020 to 53.7 percent in February 2021, suggesting a 7-day incidence rate of 2,223 per 100,000 inhabitants. In Jimma the numbers increased from 30.8 percent in November 2020 to 56.1 percent in February 2021, suggesting an incidence rate of 3,810 per 100,000 inhabitants. Urban communities showed a significant increase in seroprevalence to almost 40 percent in early 2021, suggesting incidence rates of 1,622 in Jimma and 4,646 per 100,000 inhabitants in Addis Ababa, respectively. Seroprevalence in rural areas increased from 18 percent in November 2020 to 31 percent in March 2021, well above the WHO or Johns Hopkins University figures reported to date (of 290,000 infections in a population of 112 million – equivalent to a prevalence of 0.26 percent, as of August 2021).

Call for reorientation of the vaccination strategy for Africa:

This study illustrates the dynamics of COVID-19 infections in an African population through the spread of SARS-CoV-2 among healthcare workers, and in urban and rural Ethiopia. The results imply a large wave of infection in Ethiopia, requiring special attention with regards to healthcare burden and disease prevention. In this context, Professor Michael Hoelscher, Director of the Division of Infectious Diseases and Tropical Medicine at the LMU University Hospital, calls for an adaption of the vaccination strategy for African countries: "Our results indicate that immunity – due to passed SARS-CoV-2 infections – is much higher in the African population than assumed. This allows vaccines to be used more effectively. For example, the antibody status could be determined before vaccination and, if necessary, only one dose could be vaccinated as a booster. In addition, this allows for a targeted use of the vaccine, especially in high-risk groups and the elderly."

*Partners in the study:

- Lead: Division of Infectious Diseases and Tropical Medicine, LMU University Hospital Munich, Germany

- German Center for Infection Research (DZIF), partner site Munich, Germany

- Jimma University (JU), Institute of Health, Jimma, Ethiopia

- St. Paul's Hospital (St. Paul's Hospital Millennium Medical College, SPHMMC), Addis Ababa, Ethiopia

- Institute for Medical Information Processing, Biometry, and Epidemiology IBE, LMU Munich, Germany

- Institute of Computational Biology, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany

- Center for Mathematics, Technische Universität München, Garching, Germany

- Faculty of Mathematics and Natural Sciences, University of Bonn, Bonn, Germany

Funding:

The study was funded by the Bavarian State Chancellery, by the German Federal Ministry of Education and Research (BMBF) (MoKoC019;01KI20271), by the EU funding program "Horizon 2020" of the European Commission (ORCHESTRA; 101016167), by the German Research Foundation (SEPAN; HA 7376/3-1), and by the Volkswagen Foundation (E2; 99 450)

contact for scientific information:

Prof. Dr. med. Michael Hoelscher

Director of the Division of Infectious Diseases and Tropical Medicine, LMU University Hospital

(idw)

Phone: +49 (0)89 4400-59801 E-Mail: hoelscher@lrz.uni-muenchen.de

Dr. med. Arne Kroidl Division of Infectious Diseases and Tropical Medicine, LMU University Hospital Phone: +49 (0)89 4400 59816 E-Mail: akroidl@lrz.uni-muenchen.de

PD Dr. med. Andreas Wieser Division of Infectious Diseases and Tropical Medicine, LMU University Hospital Max von Pettenkofer-Institute, LMU Phone: +49 (0)89 2180 78 296 E-Mail: wieser@mvp.lmu.de

Original publication:

Gudina E K*, Ali S*, Girma E, Gize A, Tegene B, Hundie G B, Sime, W T, Ambachew R, Gebreyohanns A, Bekele M, Bakuli A, Elsbernd K, Merkt S, Contento L, Hoelscher M, Hasenauer J, Wieser A*, Kroidl A*

Seroepidemiology and model-based prediction of SARS-CoV-2 in Ethiopia: longitudinal cohort study among front-line hospital workers and communities, The Lancet Global Health 2021; 9: e1517–27. https://www.thelancet.com/journals/langlo/article/PIIS2214-109X(21)00386-7/fulltext

(*shared first and last authorships)

URL for press release: https://www.lmu-klinikum.de/aktuelles/pressemitteilungen/sars-cov-2-verbreitung-in-athiopie n-stark-unterschatzt/oc78b78b67abbcb3

Attachment Spread of SARS-CoV-2 in Ethiopia is underestimated http://idw-online.de/en/attachment87737